

<p><b>CONSENT FORM</b> FOR <b>CAESAREAN SECTION</b></p>	<p>Med Rec. No: &lt;&lt;Patient Demographics:Record Number&gt;&gt;</p> <p>Surname:&lt;&lt;Patient Demographics:Surname&gt;&gt;</p> <p>Forename: &lt;&lt;Patient Demographics:First Name&gt;&gt;</p> <p>DOB:&lt;&lt;Patient Demographics:DOB&gt;&gt;</p>
<p><b>An incision is made in the abdomen either in the midline below the umbilicus, or in the hairline just above the pubic bone.</b></p> <p><b>A vertical (Classical) or horizontal (Lower Segment) incision is made on the uterus and the baby (s) and placenta are delivered.</b></p> <p><b>General, Regional or Local anaesthetic is used.</b></p>	

**RISKS**

*These are the more common risks. There may be other unusual risks that have not listed here. Please ask Dr Rana if you have any general or specific concerns.*

I understand there are risks associated with any **anaesthetic**, and I can discuss these with the Anaesthetist. I may have side effects from any of the drugs used. The more common side effects include light-headedness, nausea, skin rash and constipation.

- Wound. Pain , bruising and redness is common.
- There is a 1/300 risk of a hematoma requiring drainage. Longer term risks include Hernia formation (less than 1/1000 risk), and Keloid scar formation.
- Haemorrhage. The risks of significant bleeding is less than 5%. The risk of requiring a transfusion, or further surgery to deal with bleeding, is less than 1%.
- Infection. This may occur in various sites, including the chest, pelvis, wounds and bladder. In most cases infection can be treated with antibiotics, but on occasions may require further surgery.
- Damage to other organs. These include bowel/rectum (risk less than 2%), bladder (risk less than 2%), or the ureter ( tube from kidney to bladder) (risk less than 1%). If this happens the usual procedure is to carry out immediate repair. This may include bowel surgery, colostomy, bladder or ureteric repair. On occasions this damage may not be immediately recognised, and so there may be a delay in diagnosis, and subsequent repair.
- Bladder, bowel and sexual function. There may be short-term and long-term changes in function. Urinary catheterisation may be required.
- Venous thromboembolism. Blood clots can develop in the veins of the legs or pelvis, which may cause pain and swelling. If these become loose, they may travel to the lungs (Pulmonary embolus), making me short of breath. Occasionally this may be fatal. Patients are frequently given elasticated stockings to protect against clot formation, and some patients are given anticoagulant injections.
- Fluid may develop in the pelvis or the top of the vagina, and may require antibiotics or further surgery.
- Electric diathermy can short-circuit, used to control bleeding and cause a skin burn.
- A fistula is the rare (less than one in 5000) complication involving the development of an abnormal opening between two adjoining structures, and symptoms depend on the structures involved. Repair usually involves complex surgery.
- Scar tissue can cause adhesions between loops of bowel, and there is a small risk of subsequent bowel blockages, which may require further treatment or surgery.
- Very rarely severe complications could result in death.
- Small cuts can be made on the part of the baby under the incision on the uterus. These usually heal with minimal scarring.
- During a future pregnancy the scar on the uterus may rupture. This can cause a serious threat to the life of the mother and the unborn baby. This happens in approximately 1 in 50 pregnancies following a classical caesarean section and 1 in 200 following a lower segment caesarean section.
- I am more likely in future pregnancies to have a placenta that becomes abnormally implanted. This could lead to very heavy periods after Caesarian Section and may require a Hysterectomy.

<b>DECLARATION BY PATIENT:</b>			
<ul style="list-style-type: none"> <li>• I acknowledge that <b>Dr Rana</b> has informed me about the procedure, alternative treatments and answered my specific queries and concerns about this matter.</li> <li>• I acknowledge that I have discussed with <b>Dr Rana</b> any significant risks and complications <b>specific to my individual circumstances</b> that I have considered in deciding to have this operation.</li> <li>• I agree to <b>any other additional procedures</b> considered necessary in the judgement of my obstetrician during this operation.</li> <li>• I consent to a <b>blood transfusion</b>, if needed.</li> <li>• I agree to the disposal by the hospital authorities of any tissues or parts that may be removed during the operation. I understand that some tissues or samples may be kept as part of my hospital records.</li> <li>• I have received a copy of this form to take home with me.</li> <li>• If any staff member is exposed to my blood (needle stick injury) then I consent to a sample of my blood being collected for testing for infectious diseases, such as hepatitis B, C and HIV. I understand that the blood sample will not be tested until I have been informed and I have given my consent.</li> </ul>			
<b>Signature of Patient</b>		<b>Date</b>	
<b><i>If the patient is unable to give consent, a proxy form must be completed and attached.</i></b>			
<b>DECLARATION BY DOCTOR:</b>			
<ul style="list-style-type: none"> <li>• I declare that I have explained the nature of the patient's condition, the procedure to be performed, and discussed the risks that particularly concern the patient.</li> <li>• I have given the patient an opportunity to ask questions and I have answered these.</li> </ul>			
<b>Doctor's Signature</b>		<b>Date</b>	
<b>Doctor's Name</b> <i>(please print)</i>	Dr Ritu Rana		
<b>INTERPRETER'S DECLARATION:</b>			
I confirm I have accurately interpreted the contents of this form and the related conversations between the patient and the doctor.			
<b>Interpreter's Signature</b>		<b>Date</b>	
<b>Interpreter's Name</b> <i>(please print)</i>			